

alcohol and other drug use in young people

for family and friends



Adolescence and young adulthood is an exciting time of change in all aspects of life: the body, thoughts, feelings and relationships with others.

Some of the brain changes that happen during adolescence lead young people to do things that feel good or exciting. At the same time, there are areas of the brain that are still developing. These areas have a role in being able to plan ahead, manage emotions and resist peer pressure.¹ This means that young people are more likely to try out new experiences and take risks than children or older adults, especially when they have strong feelings,² or when they are with friends.³

Taking risks is a normal part of growing up. It can be part of a young person forming their own identity, practicing making decisions, developing new skills, learning the consequences of their actions and becoming more independent.⁴ However risk-taking can sometimes get people into unsafe situations too.



About this resource

This resource is for family or friends of young people aged 12-25 years. It aims to:

- answer common questions about alcohol and other drugs (AOD)
- provide practical information about what you can do if you are worried about your young person using AOD, and
- give guidance on supporting a young person who is experiencing problems related to AOD.

For the purpose of this resource 'adolescents' refers to young people aged 12-17 and 'young adults' means those aged 18-25 years.

Alcohol and other drugs

Drugs, including alcohol, are substances that can quickly change how someone thinks, feels or behaves. Drugs can be legal (e.g., alcohol or tobacco), or illegal (e.g., cannabis or cocaine). Illicit drug use means:

- using illegal drugs
- using pharmaceuticals for non-medical purposes, like using prescription drugs without a prescription, or using them in an amount or for a reason for which they aren't intended to be used
- using certain substances inappropriately, such as sniffing petrol.⁵

Drugs are grouped by the types of effect they have on the brain. You can go to adf.org.au/drug-facts/#wheel to see the categories, their effects and examples.

Why do people use alcohol and other drugs?

People use alcohol and other drugs for many different reasons. The most common reasons people first decide to try AOD are curiosity, because friends or family offered it or were using it, or to do something exciting.⁵ Other reasons why people use AOD are:

- to relax
- for enjoyment or pleasure
- to be part of a group
- to avoid physical and/or psychological pain
- to rebel
- to cope with problems
- to feel more confident
- to manage aspects of living (i.e., to stay awake at work or for study)
- to relieve stress
- to relieve boredom.^{6,7}

The reasons for trying AOD may be different to the reasons someone continues to use.⁸



There are some things that make it more or less likely that a young person will try or use AOD, or develop problematic AOD use. They are known as risk and protective factors.⁹ For example, growing up around family who are heavy drinkers may be a risk factor, while being close to family might be a protective factor.¹⁰ Things like genetics, and factors related to family, community, school, friends and life experiences all play a role.^{9, 11-13} Usually it is multiple factors, and rarely a single factor that lead to problematic AOD use. Understanding and addressing the reasons why someone uses AOD can have an impact on making changes to their AOD use.¹⁴

How many young people use AOD?

It's not unusual for young people to experiment with alcohol and other drugs. Alcohol, cannabis and tobacco are the most commonly used substances among young people in Australia.⁵

Whether or not your young person has tried AOD, chances are they will have the opportunity to do so at some point.

AOD use among young people



1 in 3

adolescents have drunk alcohol in the past year.



1 in 2

young adults have tried an illicit drug at some stage.



1 in 8

adolescents have tried an illicit drug at some stage.



81%

of young adults have drunk alcohol in the past year.^{15, 16}

AOD use and mental health

A person is most likely to experience mental ill-health during adolescence and early adulthood than at other stages of their life.¹⁷ It is also the time when many people try AOD for the first time.^{18, 19} It is common for young people who have mental ill-health to also use AOD, and vice versa.²⁰⁻²² It can sometimes be difficult to tell the difference between changes in your young person because of mental ill-health or AOD use, or due to other factors like stress.

There are different ways in which AOD use and mental ill-health can be linked. AOD use can be a consequence of mental ill-health, for example a person may use AOD to 'self-medicate' or cope with the symptoms of mental-ill health.²³ Conversely, some mental health disorders may result in someone being more likely to experiment with AOD.²⁴

On the other hand, AOD use can also lead to, or worsen, mental-ill health, with the effects of AOD intoxication, withdrawal, or long-term use triggering changes in thoughts, feelings, actions and ultimately mental illness.²⁵ This does not necessarily mean that AOD use causes mental ill-health or vice versa, but each one can increase the likelihood of the other occurring, or one could make the symptoms of the other worse.

AOD use and mental ill-health may also co-occur because there may be common factors that increase the likelihood that they will occur.²⁶ For example, genetics and trauma play a role in the development of both AOD and mental health conditions.²⁷ It is important to recognise that substance use disorders are also a type of mental health disorder in their own right.

AOD use may make the symptoms of mental ill-health, like depression anxiety and psychosis worse by making it harder to manage work, relationships, health and safety.⁵ Mental ill-health can also make recovery from substance use disorders more complicated.²⁸⁻³⁰ If a young person is taking medication as part of treatment for a mental illness and is using AOD, they may impact upon each other (i.e., making the effects of the medication and/or the substance stronger or weaker) or have unexpected negative effects on the body.³¹ Some substance use disorders involve medication as part of their treatment. Therefore, it's important to talk about any AOD use with a health profession who is prescribing medication.



Considering that AOD usage and mental-ill health can be intertwined, if your young person is using AOD and is experiencing mental-ill health, it's best to find a service that can work with both, or two services that can work closely together.³² If this isn't happening it's worth you or your young person having a chat with someone involved in their care to find a way that they can work together with you all.

Negative consequences of alcohol and other drugs

Simply telling people that something is bad for them isn't an effective way to stop them from doing it.³³ Try to avoid telling your young person about the negative aspects of AOD use without having a two-way conversation with them. The conversation may be more helpful if it is focused on the 'why' of using AOD rather than educating on the negatives of AOD. However, when they are ready, this information could be part of talking with your young person about ways they can minimise their risk of harm if they do use AOD.

There is no 100% safe or 'risk-free' level of AOD use. Use of any substance always carries some risk. When it comes to alcohol, Australian guidelines recommend that to reduce the risk of injury and other harms to health, people under 18 years of age should not drink alcohol. For those aged 18 and over, it is recommended to consume no more than 10 standard drinks per week, and no more than four in one day.³⁴



AOD use poses both short-term and long-term risk of harm to young people. Some potential harms are related to drinking or using too much on one occasion, while others are related to regular use. About one third of 15-25-year-old Australians drink at 'risky' levels (e.g., more than four standard drinks on one occasion) on a monthly basis.³⁵

Short-term or immediate risks of AOD use in young people include:

- Heightened risk of experiencing an accident, injury, or death. For example, from alcohol poisoning, road traffic accidents, suicide and self harm, falls, and drowning.³⁶⁻³⁸
- Increased likelihood of risky sexual behaviour, such as unsafe sex (resulting in sexually transmitted infections or unintentional pregnancy)³⁶. Also increased risk of non-consensual sex, either as a victim or perpetrator.^{39, 40}
- Increased risk of experiencing violence such as being a victim of assault or getting into fights.³⁶

Longer-term risks are:

- Potential negative effects on the brain's structure and function. Adolescence is an important time for brain development, and it continues to develop into our mid-twenties. This means that young people's brains are particularly vulnerable to the effects of AOD.^{41, 42}
- Physical health problems including liver and cardiovascular disease, cancers, and obesity.³⁶
- Higher levels of use later in life and increased risk of AOD-related problems later in life, including developing a substance use disorder.^{43, 44} About half of the people who develop a SUD do so before the age of 20.¹⁷

For information about specific drugs and their short and long-term effects, see adf.org.au/drug-facts/#list

Signs that AOD use is problematic

The majority of people who use AOD will not experience problems relating to their use, but some people do.

Broadly, AOD use is considered to be a problem when a person has developed a substance use disorder (SUD), for example alcohol use disorder, opioid use disorder etc. A SUD is a mental illness diagnosed by a health professional. A person's AOD use needs to meet certain criteria for them to be diagnosed with a SUD. A SUD can be mild, moderate or severe. A person may have a SUD when:

- Their AOD use has frequent harmful effects on health and wellbeing. For example, using in situations that are physically risky, or continuing to use despite experiencing physical or mental health problems that are caused by the substance or made worse by it.
- They have less control over their AOD use than they would like. For example, using more of the substance/s or using over a longer time period than originally planned, not being able to cut down despite wanting or trying to, spending a lot of time getting the substance, using it and recovering from its effects.
- They experience cravings. These can happen at any time but particularly in situations where the substance was previously gotten or used.
- AOD interferes with important activities like school, work, hobbies and/or relationships with family and friends.
- They need to increase how much of the substance is used to get the same effects, or there is a reduced effect when the usual amount is used (known as 'tolerance'). The person may also develop 'withdrawal' symptoms, which means experiencing unpleasant physical or mental symptoms after using less of a substance or not having it.⁴⁵

The signs of problematic AOD use in your young person could be social, psychological or physical.

You might see:

- Changes in eating habits or weight
- Changes in sleeping patterns
- Sudden or frequent changes in mood or behaviour
- Fluctuating energy levels (hyperactive and agitated/ fatigued and sullen)
- Slurred speech or being unsteady on their feet
- Changes in performance or attendance at school, work or other activities
- Loss of interest in hobbies
- Money problems or a need for money
- Changes in friendships, whether it be new friends or seeing friends less often.

Each of these signs could also be explained by something else such as stress, mental health problems or just normal changes as part of adolescence. If you are noticing changes in your young person, open and supportive communication is the most important way to find out what is happening for them.

How you can support your young person

Practical tips

There is no guaranteed way to prevent your young person from experimenting with AOD. People from any background use AOD or can experience problems relating to their use. However, family and friends can play an influential role when it comes to AOD use in their young person. There are also lots of things you can do that can help prevent harmful use and also to support a young person who is engaging in problematic use.⁴⁶

Role model responsible behaviour with AOD

Your opinions and behaviours towards alcohol and other drugs have an influence on your young person.⁴⁶ Demonstrate a healthy approach to AOD by doing things like:


- drinking in moderation if you drink alcohol (i.e., following the Australian guidelines – nhmrc.gov.au/health-advice/alcohol)
- showing that you can still socialise and have fun without always involving AOD
- modelling ways to relax or cope with stress that don't include AOD (e.g., meditation, exercise, talking to a friend, getting enough sleep and eating well). This includes being mindful of commonly used phrases that unintentionally role model that AOD is needed to cope in certain situations, such as "I need a drink"
- demonstrating ways in real life of saying no to AOD when you don't want to have any or you have had enough
- never driving or operating machinery while under the influence of AOD (e.g., drinking and driving)
- using medications as directed.^{47, 48}

Manage access to AOD

Avoid giving any alcohol to young people under the age of 18. Young people whose parents give them alcohol (whole drinks or a sip/taste) are more likely to start drinking at an earlier age, drink at risky levels later in life, and experience alcohol-related harms (e.g., getting sick from drinking, being in a fight, damaging property, getting in trouble with friends, parents, teachers or the police).^{46, 49} Providing alcohol can be seen by children and adolescents as 'approval' and contributes to family examples of drinking.^{50, 51} Similarly, make sure that prescription medications are stored securely, used only as directed and only by the person that they are prescribed for.

Stay connected

Having a good relationship with your young person is protective against drinking.^{46, 52} Building a good relationship involves doing activities together, whether it be hobbies, chores, watching TV or eating dinner together regularly. Encourage your young person to communicate with you about where they are, what they are doing and who they're with.⁴⁶ This is not the same as strict control or harsh discipline, which does not protect against AOD use.⁵³



Support opportunities for belonging

Having a sense of belonging or 'being part' of a school or community is protective against AOD use.^{11, 12}

This means the young person having the opportunity to get involved in activities they enjoy at school or within the community, like sports or organised groups.

More information about different influences on young people's AOD use can be found at: adf.org.au/talking-about-drugs/parenting/influencers/

Starting a conversation about alcohol and other drugs

Some things are tricky to talk about, including AOD. It can be helpful to approach these conversations as a two-way street, where you listen, hear and validate your young person's experience. Creating an environment where your young person feels safe approaching you is a protective factor against AOD use.⁴⁶ If you haven't talked about AOD with your young person before, it's not too late to start.

Before you raise the issue of AOD with your young person, it can be helpful to get some support and advice specific to your circumstances to help you feel calm and prepared to talk about your concerns. See the 'resources for family and friends' section.

Get the facts

It can be helpful to have the right information about alcohol and other drugs. There is a lot of misinformation and opinions about AOD, but if you can focus on the facts the conversation will be more successful. You can find some reliable sources in the 'helpful resources' section.

Raising your concerns

Organise a time to talk in a private place where you both feel comfortable. Sitting directly across from your young person may feel confronting or too formal for some people. Sometimes it's easier to talk while doing something side-by-side, like sitting on the couch, walking the dog or driving in the car together. Ensure you have enough time. It's also best to have conversations when your young person is not affected by AOD (or experiencing comedown/withdrawal) or if they are tired or stressed.

Begin with general and open-ended questions about how things are going (i.e., questions that have an answer longer than 'yes' or 'no'). An example could be "I've noticed you don't seem like yourself lately and I'm worried. What's going on for you right now?"

If you do ask directly about using AOD, try not to make assumptions or accusations. Give specific examples about things you are concerned about without blaming, such as, "I've noticed you're missing school/work more often and I'm concerned about you, is there anything you want to talk about, or something I could do to support you?" It's also OK to be honest and say "I'm worried that you might be drinking too much but I could be wrong".

Be prepared that your young person may not have all the answers or be able to explain. They may not want to talk to you about it at all. This can be due to lots of different things; they might not know why they use or find it hard to put things into words. They could also be afraid of your response or of getting others into trouble.

When your young person isn't worried about their AOD use or doesn't want to seek help

It is common for young people to not be concerned about their AOD use, or to not want to stop or cut back.⁵⁴ This can be difficult for those around them. Young people and their family and friends are often not on the same page about whether or not AOD use is something to be concerned about. There is also a lot of stigma connected with substance use disorders.

People can be at different stages of readiness for any type of behaviour change.⁵⁵ If they aren't ready, trying to push your young person to seek help or convince them to change may not be a good idea. It could make them feel pressured and pull away, or be more secretive. It could also make them resist reducing or stopping their AOD use. However, it's also important to think about the individual circumstances of your young person if you have noticed problematic AOD use; for example, if they are a younger adolescent, their pattern of use is more severe or there are safety concerns related to AOD use (e.g., risk of overdose, driving under the influence, self-harm or suicide). It's OK to voice your concerns. If you're unsure of what to do, reach out to one of the resources for family and friends (see below), your GP or a mental health clinician if your young person is linked in with one.

Try to talk with your young person about how they would like things to be different and what they want for the future and to reflect on AOD use in this context. Having someone think about their own reasons for making changes and coming to the decision themselves is more likely to make them want to take the next step. Let your young person know you will be there when they are ready. The good news is that health professionals have different techniques for working with young people at different levels of readiness and with different goals for AOD use.



Try to stay calm and listen to your young person

It's important to try and stay calm when talking about difficult things, including AOD. If the conversation gets heated or turns into a lecture, your young person may get defensive or shut down and refuse to talk. Try to come from a place of curiosity rather than judgement and understand why your young person has used AOD and what function it serves for them. For example, all their social events may centre around AOD use and they might worry that they won't fit in and will lose friends if they reduce or stop AOD use. Give them a chance to speak. This doesn't mean you have to agree with their actions. Having their story heard can mean your young person knows you care about them and feels they can talk to you. If things get off track it might be a good idea to pause the conversation and come back to it another time. Even if things go well it can be a continuing conversation, not everything has to be said in one go.

Navigating talking about your own experiences

Your young person may ask you about your own past experiences with AOD. Evidence suggests that parents sharing information about their own use of alcohol and/or marijuana might come across as approval of AOD, and has been linked with higher levels of use and with more negative consequences of use in their young people.^{56, 57} On the other hand, being honest may build trust and closeness between you and your young person.⁵⁶ If you do choose to talk about it, keep it brief and factual. It's not necessarily about what you did or didn't do but it can be a way to talk about why you made the choices that you did and what the consequences were.

Let them know you're always here to help

It is OK to be clear that you don't agree with AOD use, while acknowledging that your young person may choose to do it anyway. This might mean talking about ways they can keep themselves safe – things like having a plan for getting home before they go out, only taking a set number of drinks to a party or telling someone what they are taking in case anything goes wrong. Go to hrvic.org.au for ways people can lower their risk of harm if they do choose to use AOD (called 'harm reduction').

Let your young person know that you love them and that they can call you anytime if they are in trouble and that you won't be angry – their safety is the most important thing. If this isn't an option then talk about who they could call if they needed help.

Set boundaries

Family and friends can worry that setting boundaries will make things worse, but there are ways to do so while maintaining your relationship with your young person. It's a good idea to get your own support to help you to do this. You might want to work on setting boundaries around things like using AOD in your home, who comes to the house, or ways to protect other young people living with you. You'll also need to consider the consequences of breaking the rules, and making sure that other important adults in the young person's life are on the same page. A good place to start is fds.org.au/setting-boundaries.

Encourage help-seeking

Support your young person to reach out if they need to, even if they don't feel like they are able to talk to you.

Some more tips for talking to your young person about AOD use can be found at positivechoices.org.au/parents/starting-the-conversation-about-drug-use

Getting support for AOD use when you're worried

Learning that your young person has tried or uses AOD does not mean they have a substance use problem or need treatment. Many people can stop on their own, or with support from their family, friends or doctor. However, if you are worried that their AOD use is problematic, or your young person feels like their use is out of their control or they want to work on having a healthier relationship with AOD, it's a good idea to reach out early.

The first step is to contact your local headspace centre or ehheadspace.org.au, General Practitioner (GP), or your state's alcohol and drug information service. Go to adf.org.au/resources/help-support-state or call 1300 85 85 84 to find out what is available locally and receive free and confidential advice. If your young person is already engaged in mental health care, speak to their mental clinician about your concerns with your young person's consent, or discuss with your young person speaking to their mental health worker.

There are specialist AOD treatment services that work to help people with problems related to AOD use. The goal is to support the person to reduce or stop AOD use, or to work with them to reduce harm to themselves, their family and the community when they do use AOD.

There are many different options available to support people to change their AOD use, including some specifically for young people. See adf.org.au/talking-about-drugs/seeking-help/ for more information. Many specialist AOD services do not need a doctor's referral.

Treating problematic AOD use is not as simple as just stopping. Completely stopping AOD use suddenly often isn't realistic or safe.

Confidentiality

You or your young person may want to reach out for help but be worried about the potential consequences, like school finding out or the police being informed. It is against the law for health services to share information without the client's permission, unless they are worried about someone's immediate safety. This should be explained in the first meeting with a health service.

Looking after yourself

Supporting a young person who experiences problems with AOD can be challenging. When travelling on a plane, in the event of an emergency passengers are advised to put on their own oxygen mask first so that they can assist a fellow passenger. Looking after yourself is an important part of being able to support others. For more information and ideas about self-care go to: headspace.org.au/friends-and-family/self-care-for-family-and-friends

There are services especially for family and friends of people who use AOD. More information is on the next page of this resource.

Sometimes people can use threatening, abusive or violent behaviour when under the influence of AOD, or as a way to get others to help them access AOD. This could be physical violence, property damage, verbal abuse, controlling behaviour or financial abuse. Violence is never OK. Contact 1800 RESPECT or 1800respect.org.au for confidential information and support. If someone is at immediate risk of harm to themselves or others, call 000.



Resources for family and friends

Family Drug Support: Australia-wide support for families and friends of people using drugs and alcohol, including a 24/7 telephone support line, support groups, education programs and counselling for families. fds.org.au

Counselling Online: A free, confidential, 24/7 online text-based counselling service for people across Australia concerned about their drinking or drug use, or the use of a family member or friend. counsellingonline.org.au

Family and Friend Support Program: a free online program designed for people supporting a loved one who uses alcohol and/or other drugs. aod.ffsp.com.au

eheadspace: eheadspace provides free online and telephone support and counselling to young people 12 – 25 and their families and friends. headspace.org.au/eheadspace

Helpful resources

Alcohol and Drug Foundation adf.org.au

Your Room yourroom.health.nsw.gov.au/Pages/home.aspx

Australian Government health.gov.au/health-topics/drugs and health.gov.au/health-topics/alcohol

Culturally specific services

Australian Indigenous Health Infonet: alcohol and other drugs knowledge centre aodknowledgecentre.ecu.edu.au

Written resources in a number of languages other than English, select 'filter by > languages' yourroom.health.nsw.gov.au/resources/publications/pages/publications.aspx

Authors

Alicia Randell

Orygen, Centre for Youth Mental Health, University of Melbourne

Expert reviewers

Allie Bailey

Orygen

Dr Cali Bartholomeusz

Orygen, Centre for Youth Mental Health, University of Melbourne

Desirée Smith

Orygen

Dr Eddie Mullen

Orygen, Melbourne Health, The University of Melbourne and the National Centre for Clinical Research on Emerging Drugs, The University of New South Wales

Dr Enrico Cementon

Orygen, University of Melbourne

Associate Professor Gill Bedi

Orygen, Centre for Youth Mental Health, University of Melbourne

Helen Nicoll

Orygen

Dr Steve Leicester

eheadspace

Susan Preece

Orygen

Youth Consultant

Emily Unity

Orygen

We would like to recognise individuals who responded to our online survey which informed the development of this resource.

references

- Steinberg L. A dual systems model of adolescent risk-taking. *Dev Psychobiol.* 2010;52(3):216-24.
- Figner B, Mackinlay RJ, Wilkening F, Weber EU. Affective and deliberative processes in risky choice: age differences in risk taking in the Columbia Card Task. *J Exp Psychol: Learn, Mem Cogn.* 2009;35(3):709-30.
- Gardner M, Steinberg L. Peer influence on risk taking, risk preference, and risky decision making in adolescence and adulthood: an experimental study. *Dev Psychol.* 2005;41(4):625-35.
- Kelley A, Schochet T, Landry C. Risk taking and novelty seeking in adolescence: introduction to part I. *Ann N Y Acad Sci.* 2004;1021:27-32.
- Australian Institute of Health and Welfare. National Drug Strategy Household Survey 2019. Canberra: AIHW; 2020. Contract No.: Drug Statistics series no. 32. PHE 270. Available from: <https://www.aihw.gov.au/getmedia/3564474e-f7ad-461c-b918-7f8de03d1294/aihw-phe-270-NDSHS-2019.pdf.aspx?inline=true>.
- Ritter A, King T, Hamilton MA. Drug use in Australian Society. South Melbourne, Victoria: Oxford University Press; 2013.
- Boys A, Marsden J, Strang J. Understanding reasons for drug use amongst young people: a functional perspective. *Health Educ Res.* 2001;16(4):457-69.
- Titus JC, Godley SH, White MK. A post-treatment examination of adolescents' reasons for starting, quitting, and continuing the use of drugs and alcohol. *J Child Adolesc Subst Abuse.* 2007;16(2):31-49.
- Spooner C, Hetherington K. Social determinants of drug use Sydney: National Drug and Alcohol Research Centre; 2005. Contract No.: NDARC Technical Report No.228. Available from: <https://ndarc.med.unsw.edu.au/sites/default/files/ndarc/resources/TR.228.pdf>.
- Stone AL, Becker LG, Huber AM, Catalano RF. Review of risk and protective factors of substance use and problem use in emerging adulthood. *Addict Behav.* 2012;37(7):747-75.
- Hemphill SA, Heerde JA, Herrenkohl TI, Patton GC, Toumbourou JW, Catalano RF. Risk and protective factors for adolescent substance use in Washington State, the United States and Victoria, Australia: a longitudinal study. *J Adolesc Health.* 2011;49(3):312-20.
- Beyers JM, Toumbourou JW, Catalano RF, Arthur MW, Hawkins JD. A cross-national comparison of risk and protective factors for adolescent substance use: the United States and Australia. *J Adolesc Health.* 2004;35(1):3-16.
- Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention. *Psychol Bull.* 1992;112(1):64-105.
- Dow SJ, Kelly JF. Listening to youth: adolescents' reasons for substance use as a unique predictor of treatment response and outcome. *Psychol Addict Behav.* 2013;27(4):1122-31.
- Australian Institute of Health and Welfare. Data tables: National Drug Strategy Household Survey 2019 – 4 Illicit use of drugs supplementary tables. Table 4.4: Lifetime(a) illicit use of drugs, by age and sex, 2001 to 2019 (per cent) [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/getmedia/54f66117-e846-4de0-a874-e5f5eee57214/aihw-phe-270-4-Illicit-use-of-drugs-tables.xlsx.aspx>
- Australian Institute of Health and Welfare. Data tables: National Drug Strategy Household Survey 2019 – 3 Alcohol supplementary tables. Table 3.4: Alcohol drinking status, by age and sex, 2019 (row per cent) [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/getmedia/4f178aed-4301-4d49-8fe6-c9fa663d914e/aihw-phe-270-3-Alcohol-tables.xlsx.aspx>
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005;62(6):593-602.
- Australian Institute of Health and Welfare. Data tables: National Drug Strategy Household Survey 2019 – 4 Illicit use of drugs supplementary tables. Table 4.17: Age of initiation(a) of lifetime(b) illicit use of drugs, people aged 14 and over, 2001 to 2019 (years) [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/getmedia/54f66117-e846-4de0-a874-e5f5eee57214/aihw-phe-270-4-Illicit-use-of-drugs-tables.xlsx.aspx>
- Australian Institute of Health and Welfare. Data tables: National Drug Strategy Household Survey 2019 – 3 Alcohol supplementary tables. Table 3.31: Age of initiation, recent drinkers(a) and ex-drinkers(b) aged 14 and over, 2001 to 2019 (years) [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/getmedia/4f178aed-4301-4d49-8fe6-c9fa663d914e/aihw-phe-270-3-Alcohol-tables.xlsx.aspx>
- Chan Y-F, Dennis ML, Funk RR. Prevalence and comorbidity of major internalizing and externalizing problems among adolescents and adults presenting to substance abuse treatment. *J Subst Abuse Treat.* 2008;34(1):14-24.
- Armstrong TD, Costello EJ. Community studies on adolescent substance use, abuse, or dependence and psychiatric comorbidity. *J Consult Clin Psychol.* 2002;70(6):1224-39.
- Lawrence D, Johnson S, Hafekost J, Boterhoven de Haan K, Sawyer M, Ainsley J, et al. The mental health of children and adolescents: report on the second Australian child and adolescent survey of mental health and wellbeing. Canberra: Department of Health; 2015. Available from: <https://www.health.gov.au/resources/publications/the-mental-health-of-children-and-adolescents>.
- Robinson J, Sareen J, Cox BJ, Bolton JM. Role of self-medication in the development of comorbid anxiety and substance use disorders: a longitudinal investigation. *Arch Gen Psychiatry.* 2011;68(8):800-7.
- Swann AC. The strong relationship between bipolar and substance-use disorder. *Ann N Y Acad Sci.* 2010;1187(1):276-93.
- Fiorentini A, Volonteri L, Dragogna F, Rovera C, Maffini M, Mauri M, et al. Substance-induced psychoses: a critical review of the literature. *Curr Drug Abuse Rev.* 2011;4(4):228-40.
- Marel M, Mills K, Kingston R, Gournay K, Deady M, Kay-Lambkin F, et al. Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings (2nd edition). Sydney, Australia: Centre of Research Excellence in Mental Health and Substance Use, National Drug and Alcohol Research Centre, University of New South Wales; 2016.
- Cerdá M, Sagdeo A, Johnson J, Galea S. Genetic and environmental influences on psychiatric comorbidity: a systematic review. *J Affect Disord.* 2010;126(1-2):14-38.
- Baker KD, Lubman DI, Cosgrave EM, Killackey EJ, Pan Yuen H, Hides L, et al. Impact of co-occurring substance use on 6 month outcomes for young people seeking mental health treatment. *Aust N Z J Psychiatry.* 2007;41(11):896-902.
- Grella CE, Hser Y-I, Joshi V, Rounds-Bryant J. Drug treatment outcomes for adolescents with comorbid mental and substance use disorders. *J Nerv Ment Dis.* 2001;189(6):384-92.
- Lubman DI, Allen NB, Rogers N, Cementon E, Bonomo Y. The impact of co-occurring mood and anxiety disorders among substance-abusing youth. *J Affect Disord.* 2007;103(1-3):105-12.

alcohol and other drug use in young people for family and friends

31. Kaminer Y, Goldberg P, Connor DF. Psychotropic medications and substances of abuse interactions in youth. *Subst Abus.* 2010;31(1):53-7.
32. Baker D, Kay-Lambkin F. Two at a time: alcohol and other drug use by young people with a mental illness. Melbourne: Orygen; 2016. Available from: https://www.orygen.org.au/Policy/Policy-Reports/Alcohol-and-other-drug-use/alcohol_and_other_drug_policy_paper_2016?ext.
33. Pan W, Bai H. A multivariate approach to a meta-analytic review of the effectiveness of the DARE program. *Int J Environ Res Public Health.* 2009;6(1):267-77.
34. National Health and Medical Research Council. Australian Guidelines to Reduce Health Risks from Drinking Alcohol. Canberra: NHMRC; 2020. Available from: <https://www.nhmrc.gov.au/about-us/publications/australian-guidelines-reduce-health-risks-drinking-alcohol>.
35. Australian Institute of Health and Welfare. Data tables: National Drug Strategy Household Survey 2019 – 3 Alcohol supplementary tables. Table 3.17: People at risk of injury on a single occasion of drinking, by age and sex, 2007 to 2019 (row per cent) [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/getmedia/4f178aed-4301-4d49-8fe6-c9fa663d914e/aihw-phe-270-3-Alcohol-tables.xlsx.aspx>
36. Hall WD, Patton G, Stockings E, Weier M, Lynskey M, Morley KI, et al. Why young people's substance use matters for global health. *Lancet Psychiatry.* 2016;3(3):265-79.
37. Australian Institute of Health and Welfare. Deaths web report, supplementary data table S3.2: leading causes of death, number and crude rates (deaths per 100,000 population) by age group, 2016-2018 [data file]. AIHW: Canberra 2020. Available from: <https://www.aihw.gov.au/reports/life-expectancy-death/deaths/data>
38. Australian Institute of Health and Welfare. Impact of alcohol and illicit drug use on the burden of disease and injury in Australia: Australian Burden of Disease Study 2011. Canberra: Australian Government; 2018. Available from: <https://www.aihw.gov.au/getmedia/34569d3a-e8f6-4c20-aa6d-e1554401ff24/aihw-bod-19.pdf.aspx?inline=true>.
39. Testa M. The role of substance use in male-to-female physical and sexual violence: a brief review and recommendations for future research. *J Interpers Violence.* 2004;19(12):1494-505.
40. Abbey A, Zawacki T, Buck PO, Clinton AM, McAuslan P. Sexual assault and alcohol consumption: what do we know about their relationship and what types of research are still needed? *Aggress Violent Behav.* 2004;9(3):271-303.
41. Spear LP. Effects of adolescent alcohol consumption on the brain and behaviour. *Nat Rev Neurosci.* 2018;19(4):197-214.
42. Squeglia LM, Jacobus J, Tapert SF. The influence of substance use on adolescent brain development. *Clin EEG Neurosci.* 2009;40(1):31-8.
43. Winters KC, Lee C-YS. Likelihood of developing an alcohol and cannabis use disorder during youth: association with recent use and age. *Drug Alcohol Depend.* 2008;92(1-3):239-47.
44. Magid V, Moreland AD. The role of substance use initiation in adolescent development of subsequent substance-related problems. *J Child Adolesc Subst Abuse.* 2014;23(2):78-86.
45. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Arlington, VA: American Psychiatric Association; 2013.
46. Ryan SM, Jorm AF, Lubman DI. Parenting factors associated with reduced adolescent alcohol use: a systematic review of longitudinal studies. *Aust N Z J Psychiatry.* 2010;44(9):774-83.
47. Your Room – Drug and Alcohol information. Talking about it. [Internet]: NSW Health and St Vincent's Alcohol and Drug Information Service; 2021 [cited 2021 June 1]. Available from: <https://yourroom.health.nsw.gov.au/Families/Pages/talking.aspx>.
48. Alcohol and Drug Foundation. Talking about drugs. North Melbourne: Australian Drug Foundation; 2020 [updated 17 Sept 2020; cited 2021 1 June]. Available from: <https://adf.org.au/talking-about-drugs/>.
49. Aiken A, Clare PJ, Boland VC, Degenhardt L, Yuen WS, Hutchinson D, et al. Parental supply of sips and whole drinks of alcohol to adolescents and associations with binge drinking and alcohol-related harms: a prospective cohort study. *Drug Alcohol Depend.* 2020;215:108204.
50. Jackson KM, Barnett NP, Colby SM, Rogers ML. The prospective association between sipping alcohol by the sixth grade and later substance use. *J Stud Alcohol Drugs.* 2015;76(2):212-21.
51. Donovan JE, Molina BS. Children's introduction to alcohol use: sips and tastes. *Alcohol Clin Exp Res.* 2008;32(1):108-19.
52. Yap MB, Cheong TW, Zaravinos-Tsakos F, Lubman DI, Jorm AF. Modifiable parenting factors associated with adolescent alcohol misuse: a systematic review and meta-analysis of longitudinal studies. *Addiction.* 2017;112(7):1142-62.
53. Becoña E, Martínez Ú, Calafat A, Juan M, Fernández-Hermida JR, Secades-Villa R. Parental styles and drug use: a review. *Drugs Educ Prev Policy.* 2012;19(1):1-10.
54. Reavley NJ, Cvetkovski S, Jorm AF, Lubman DI. Help-seeking for substance use, anxiety and affective disorders among young people: results from the 2007 Australian National Survey of Mental Health and Wellbeing. *Aust N Z J Psychiatry.* 2010;44(8):729-35.
55. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol.* 1983;51(3):390-95.
56. Napper LE, Derby AR. Longitudinal associations between maternal disclosure of past alcohol use and underage college drinking. *Psychol Addict Behav.* 2018;32(3):301-8.
57. Napper LE, Froidevaux NM, LaBrie JW. Being blunt about marijuana: parent communication about marijuana with their emerging adult children. *Prev Sci.* 2016;17(7):882-91.



headspace would like to acknowledge Aboriginal and Torres Strait Islander peoples as Australia's First People and Traditional Custodians. We value their cultures, identities, and continuing connection to country, waters, kin and community. We pay our respects to Elders past and present and are committed to making a positive contribution to the wellbeing of Aboriginal and Torres Strait Islander young people, by providing services that are welcoming, safe, culturally appropriate and inclusive.



headspace is committed to embracing diversity and eliminating all forms of discrimination in the provision of health services. headspace welcomes all people irrespective of ethnicity, lifestyle choice, faith, sexual orientation and gender identity.



headspace centres and services operate across Australia, in metro, regional and rural areas, supporting young Australians and their families to be mentally healthy and engaged in their communities.

**For more details about headspace
visit headspace.org.au**

headspace National Office

p +61 3 9027 0100

f +61 3 9027 0199

info@headspace.org.au

Acknowledgements

This resource was prepared by the Centre of Excellence in Youth Mental Health. The series aims to answer common questions using the research evidence about mental health and substance use problems affecting young people. Experts on the topic have reviewed the summary before publication. The authors would like to thank the clinical consultants and youth contributors for their input on this resource.

Disclaimer

This information is not medical advice. It is generic and does not take into account your personal circumstances, physical wellbeing, mental status or mental requirements. Do not use this information to treat or diagnose your own or another person's medical condition and never ignore medical advice or delay seeking it because of something in this information. Any medical questions should be referred to a qualified healthcare professional. If in doubt, please always seek medical advice.